

MALONATE BROTH

Medium for enterobacteria differentiation.

TYPICAL FORMULA(g/l)Ammonium Sulfate2.0Dipotassium Phosphate0.6Monopotassium Phosphate0.4Sodium Chloride2.0Sodium Malonate3.0Brom Thymol Blue0.025Final pH = 6.7 ± 0.1 at $25 \, ^{\circ}$ C.

DIRECTIONS

Suspend 8.0 g of powder in 1 liter of distilled or deionized water. Heat until completely dissolved. Dispense into final tubes. Sterilize in autoclave at 121 °C for 15 minutes. Avoid introducing extraneous carbon and nitrogen.

DESCRIPTION

MALONATE BROTH is used for differentiating Enterobacter from Escherichia based on malonate utilization.

TECHNIQUE

Inoculate tubes with a loopful of test organisms. Incubate at 36 ± 1 °C for 18-48 hours examine tubes for a change in the color of the medium from green to blue that is the sign of malonate utilization.

QUALITY CONTROL

Dehydrated medium

Appearance: free flowing, homogeneous.

Color: light green.

Prepared medium

Appearance: clear.

Color: green.

Incubation conditions: 36 ± 1 °C for 18-48 hours.

| Microorganism | ATCC | Growth | Color |
|------------------------|-------|--------------|-------|
| Escherichia coli | 25922 | poor to fair | green |
| Klebsiella pneumoniae | 13883 | good | blue |
| Salmonella typhimurium | 14028 | none | |
| Enterobacter aerogenes | 13048 | good | blue |

STORAGE

The powder is very hygroscopic: store the powder below 30 °C, in a dry environment, in its original container tightly closed and use it before the expiry date on the label or until signs of deterioration or contamination are evident. Store prepared tubes at 2-8 °C.

REFERENCES

- 1. Leifson, E. (1933). J. Bacteriol. 26: 329.
- 2. Bacteriological Analytical Manual. (1995). 8th AOAC International. Gaithersburg, MD.

PRESENTATION

| Product | REF | Σ |
|-------------------------|--------|----------|
| MALONATE BROTH (62.5 I) | 610172 | 500 g |
| MALONATE BROTH (12.5 I) | 620172 | 100 g |

TABLE OF SYMBOLS

| LOT Batch code | Caution, consult accompanying documents | Manufacturer | Contains sufficient for <n> tests</n> | IVD In Vitro Diagnostic Medical Device |
|----------------------|---|--------------|---------------------------------------|--|
| REF Catalogue number | Fragile, handle with care | Use by | Temperature limitation | Keep away from heat source |



LIOFILCHEM s.r.l.

Via Scozia, Zona Ind.le - 64026, Roseto D.A. (TE) - ITALY Phone +390858930745 Fax +390858930330





